

Mechanisms of Development 54 (1995) 257-258



Volume contents

Volume 54 (1996)

Vol. 54 No. 1		C. Chazaud, M. Oulad-Abdelghani, P. Bouillet, D. Décimo, P. Chambon, P. Dollé	
Issue contents	1	AP-2.2, a novel gene related to AP-2, is expressed in the forebrain, limbs and face during mouse embryo-	
Editorial	3	genesis	83
Introducing new editors	5	H.L. Hellmich, L. Kos, E.S. Cho, K.A. Mahon, A.	
Prof. Alexandra L. Joyner		Zimmer	
Prof. Claudio D. Stern		Embryonic expression of glial cell-line derived neu- rotrophic factor (GDNF) suggests multiple develop-	
ISDB News	7	mental roles in neural differentiation and epithelial- mesenchymal interactions	95
F. van der Hoeven, P. Sordino, N. Fraudeau, JC, Iz-			
pisúa-Belmonte, D. Duboule		M. Polimeni, S. Giorgi, L. De Gregorio, T.A. Dragani,	
Teleost HoxD and HoxA genes: comparison with		M. Molinaro, G. Cossu, M. Bouché	
tetrapods and functional evolution of the HOXD complex	9	Differentiation dependent expression in muscle cells of ZT3, a novel zinc finger factor differentially expressed in embryonic and adult tissues	107
F. Aberger, G. Schmidt, K. Richter			
The Xenopus homologue of hepatocyte growth factor-		R.J. Diaz, R. Harbecke, J.B. Singer, F. Pignoni, W.	
like protein is specifically expressed in the presump-		Janning, J.A. Lengyel	
tive neural plate during gastrulation	23	Graded effect of <i>tailless</i> on posterior gut development: molecular basis of an allelic series of a nuclear	
A. Vaahtokari, T. Åberg, J. Jernvall, S. Keränen, I. Thesleff		receptor gene	119
The enamel knot as a signaling center in the develop-			
ing mouse tooth	39	Vol. 54 No. 2	
B. Holewa, E. Pogge v. Strandmann, D. Zapp, P. Lorenz, G.U. Ryffel		Issue contents	131
Transcriptional hierarchy in <i>Xenopus</i> embryogenesis: HNF4 a maternal factor involved in the developmental		R. Schnabel, C. Weigner, H. Hutter, R. Feichtinger, H. Schnabel	
activation of the gene encoding the tissue specific transcription factor $HNF1\alpha$ (LFB1)	45	mex-1 and the general partitioning of cell fate in the early C. elegans embryo	133
	10		100
K. Ohta, M. Nakamura, K. Hirokawa, S. Tanaka, A.		A. von Bubnoff, J.E. Schmidt, D. Kimelman	
Iwama, T. Suda, M. Ando, H. Tanaka		The Xenopus laevis homeobox gene Xgbx-2 is an	
The receptor tyrosine kinase, Cek8, is transiently ex- pressed on subtypes of motoneurons in the spinal cord		early marker of anteroposterior patterning in the ecto- derm	149
during development	59		
		K. Tashiro, O. Tooi, H. Nakamura, C. Koga, Y. Ito, H.	
M. Kühl, S. Finnemann, O. Binder, D. Wedlich		Hikasa, K. Shiokawa	
Dominant negative expression of a cytoplasmically		Cloning and expression studies of cDNA for a novel	
deleted mutant of XB/U-cadherin disturbs mesoderm	7.	Xenopus cadherin (XmN-cadherin), expressed mater-	161
migration during gastrulation in Xenopus laevis	71	nally and later neural-specifically in embryogenesis	161

R. Ohlsson, A. Berkenstam

bryogenesis

Expression of a novel member of estrogen response element-binding nuclear receptors is restricted to the early stages of chorion formation during mouse em-

J. van der Wees, P.J. Matharu, K. de Roos, O.H.J. Destrée, S.F. Godsave, A.J. Durston, G.E. Sweeney Developmental expression and differential regulation by retinoic acid of <i>Xenopus COUP-TF-A</i> and <i>COUP-TF-B</i>	173	 C.J.M. de Vries, J. de Boer, J. Joore, U. Strähle, T.A.E. van Achterberg, D. Huylebroeck, K. Verschueren, K. Miyazono, A.J.M. van den Eijnden-van Raaij, D. Zivkovic Active complex formation of type I and type II activin and TGFβ receptors in vivo as studied by overexpres- 	
R. Zwart, L. Broos, G. Grosveld, D. Meijer		sion in zebrafish embryos	225
The restricted expression pattern of the POU factor		, , , , , , , , , , , , , , , , , , , ,	
Oct-6 during early development of the mouse nervous		F.J. Casado, C. Pouponnot, JC. Jeanny, O. Lecoq,	
system	185	G. Calothy, A. Pierani QR1, a retina-specific gene, encodes an extracellular	
J.B. Sass, E.S. Weinberg, P.H. Krone Specific localization of zebrafish hsp90α mRNA to myoD-expressing cells suggests a role for hsp90α		matrix protein exclusively expressed during neural retina differentiation	237
during normal muscle development	195	Author index	251
		Subject index	253
S. Staab, M. Steinmann-Zwicky Female germ cells of <i>Drosophila</i> require zygotic <i>ovo</i> and <i>otu</i> product for survival in larvae and pupae re-		Volume contents	257
spectively	205		
K. Pettersson, K. Svensson, R. Mattsson, B. Carlsson,			

211